

Volume 29, Issue 1

A generalization of monotone comparative statics: Correction

Koji Shirai Graduate School of Economics, Waseda University

Abstract

We have made a correction to "A generalization of monotone comparative statics", which is published in Economics Bulletin Vol. 3, No. 39. We correct the following three aspects of the original paper: the first and the second are the name and the definition of some fundamental notions, respectively. The third is the proof of the main proposition [Proposition 2.1, pp.5].

1 Correction

We have made a correction to "A generalization of monotone comparative statics", which is published in *Economics Bulletin Vol.3*, *No.39*. In that paper, we generalized Milgrom and Shannon's Theorem (Milgrom and Shannon (1994)) from a partially ordered set to a preordered set. As a result, we showed the following two necessary and sufficient relations. The first is the equivalence of the "w-quasisupermodularity" of objective function and the monotonicity of the solution of the constrained optimization problem with respect to "w-strong set order". The second is the similar relation of the "s-quasisupermodularity" of the function and the monotonicity of solution with respect to "s-strong set order".

We correct the following three aspects of the original paper: The first is the name of a fundamental notion. In our main proposition, the notion that is called "prelattice" in the original paper plays a crucial role. However, it turns out that the term "prelattice" has already been used as the name of a mathematical notion that is different from ours. Hence, we alter the name of ours to "preordered lattice structure". The second is the definition of s-quasisupermodularity. Thanks to a private communication from Richard Ruble, it has become clear that the original version of s-quasisupermodularity is not the necessary and sufficient condition of the monotonicity of the solution of the optimization problem with respect to the s-strong set order but only a necessary condition of it. Hence, we alter the definition of squasisupermodular in such a way that the necessary and sufficient relation is realized. The third is the proof of [Proposition 2.1, pp.5] in the original paper, specifically, the necessity part of it. It turns out that the original version is incomplete. Hence, we intend to replace it. For these purposes, we introduce some basic notions as follows.

Definition 1: Let X be a preordered set endowed with a preorder \leq . We say that $U_{x,y}$ is the set of upper bounds of $x, y \in X$ if $x \leq u$ and $y \leq u$ for all $u \in U_{x,y}$. Similarly, we say that $L_{x,y}$ is the set of lower bounds of $x, y \in X$ if $l \leq x$ and $l \leq y$ for all $l \in L_{x,y}$.

Definition 2: We say that $A_{x,y} \subset U_{x,y}$ is the set of supremums of $x, y \in X$ if $a \leq u$ for all $a \in A_{x,y}$ and $u \in U_{x,y}$. Similarly, we say that $T_{x,y} \subset L_{x,y}$ is the set of infimums of $x, y \in X$ if $l \leq t$ for all $t \in T_{x,y}$ and $l \in L_{x,y}$.

The following is our first correction: the name of this notion is altered from the original paper. In Shirai (2008), we called this a "prelattice".

Definition 3: We say that X is a preordered lattice structure if $A_{x,y} \neq \emptyset$ and $T_{x,y} \neq \emptyset$ for every $x, y \in X$.

Then, we proceed to the second and the third corrections. We introduce the following notion and lemmas. In particular, the lemmas stated below play fundamental roles in the third correction: that is, in the corrected version of the proof of Proposition 2.1 in the original paper.

Definition 4: We say that x and y are indifferent to each other if we have both $x \leq y$ and $y \leq x$. We write this as $x \sim y$ or $y \sim x$. We define the set $I_x = \{y \in X \mid x \sim y\}$, which is called *the indifference set of* x in the rest of this paper.

Lemma 1: If $x' \in I_x$ for some x, then $U_{x,y} = U_{x',y}$ and $L_{x,y} = L_{x',y}$ for every $y \in X$.

Proof. Let $u \in U_{x,y}$ and $u' \in U_{x',y}$. By the definition, $x \preccurlyeq u$ and $y \preccurlyeq u$. Since $x \sim x'$, by transitivity, we have $x' \preccurlyeq u$ and thus $U_{x,y} \subset U_{x',y}$. By similar arguments, we can prove that $U_{x',y} \subset U_{x,y}$, hence $U_{x,y} = U_{x',y}$. The rest of our claim also follows from almost the same arguments. [Q.E.D.]

Lemma 2: If $b \sim a$ for some $a \in A_{x,y}$, then $b \in A_{x,y}$. Moreover, $A_{x,y} = I_a$ for every $a \in A_{x,y}$. Similarly, if $c \sim t$ for some $t \in T_{x,y}$, then $c \in T_{x,y}$, and we have $T_{x,y} = I_t$ for every $t \in T_{x,y}$.

Proof. By transitivity, we must have $x \leq b$, $y \leq b$ and $b \leq u$ for all $u \in U_{x,y}$. This proves that $b \in A_{x,y}$. Since it is obvious that $a \sim a'$ for all $a, a' \in A_{x,y}$, our claims on $A_{x,y}$ follow. For the proof of the claims on $T_{x,y}$, we use the same logic as above. [Q.E.D.]

Lemma 3: If $x' \in I_x$ for some x, then we have $A_{x,y} = A_{x',y}$ and $T_{x,y} = T_{x',y}$ for every $y \in X$.

Proof. Let $a \in A_{x,y}$ and $a' \in A_{x',y}$. By the previous lemma, it is sufficient to show that $a' \in I_a$. By the definition, we have $x \leq a$ and $a \leq u$ for all $u \in U_{x,y}$. By transitivity and Lemma 1, this means $a \leq x'$ and $a \leq u'$ for all $u' \in U_{x',y}$ and thus, $A_{x,y} \subset A_{x',y}$. The converse relation can be shown similarly, hence $A_{x,y} = A_{x',y}$. The proof of the claim on the set of infimums is similar. [Q.E.D.]

We introduce the following notions, which play central roles in our propo-

sition. In particular, as mentioned in the opening sentence, the definition of "s-quasisupermodularity" in Definition 6 is altered from the original paper Shirai (2008). This is our second correction.

Definition 5: Let X be a preordered lattice structure and $S, S' \subset X$. We say that S' is higher than S with respect to w-strong set order if $A_{x,y} \cap S' \neq \emptyset$ and $T_{x,y} \cap S \neq \emptyset$ for every $x, y \in X$. We write this as $S \leq_{wa} S'$. We say that S' is higher than S with respect to s-strong set order if $A_{x,y} \subset S'$ and $T_{x,y} \subset S$. We write this as $S \leq_{sa} S'$.

Definition 6: Let X be a preordered lattice structure. We say that a function $f: X \to \mathbb{R}$ is w-quasisupermodular if

$$\forall t \in T_{x,y}; f(x) \ge (>)f(t) \Rightarrow \exists a \in A_{x,y}; f(a) \ge (>)f(y).$$

We say that f is s-quasisupermodular of

$$\exists t \in T_{x,y}; f(x) \ge (>) f(t) \Rightarrow \forall a \in A_{x,y}; f(a) \ge (>) f(y).$$

It should be noted that, in the original paper, s-quasisupermodularity is defined as

$$\forall t \in T_{x,y}; f(x) \ge (>)f(t) \Rightarrow \forall a \in A_{x,y}; f(a) \ge (>)f(y).$$

However, it has become clear that this definition is too weak to assure the monotonicity of the solution set of the maximization problem with respect to s-strong set relation. In the following, we write the solution set of the maximization problem: $\max_{x \in S \subset X} f(x)$ as M(S). Our main proposition can now be stated as follows. This is nothing but the corrected version of *Proposition 2.1* in the original paper.

Proposition 1: (a): Let X be a prelattice, $S, S' \subset X$, and $S \leq_{sa} S'$. Then, $M(S) \leq_{wa} M(S')$ if and only if $f: X \to \mathbb{R}$ satisfies w-quasisupermodularity. (b): $M(S) \leq_{sa} M(S')$ if and only if f satisfies s-quasisupermodularity.

Proof. (a): For the sufficiency part, see the original paper. The necessity part can be shown as follows. Let $S = I_x \cup T_{x,y}$ and $S' = I_y \cup A_{x,y}$ for some $x, y \in X$. It is obvious that $S \leq_{sa} S'$. Assume $M(S) \leq_{wa} M(S')$. Suppose $x \in M(S)$ and $y \in M(S')$. Since $M(S) \leq_{wa} M(S')$, $A_{x,y} \cap M(S') \neq \emptyset$, which implies $f(a) \geq f(y)$ for some $a \in A_{x,y}$. Suppose $x \notin M(S)$, $f(x) \geq f(t)$ for all $t \in T_{x,y}$ and $y \in M(S')$. Then, there exists some $x' \in I_x$ and $x' \in M(S)$.

By our assumption, we have $A_{x',y} \cap M(S') \neq \emptyset$ and by Lemma 3, we must have $A_{x,y} \cap M(S') \neq \emptyset$. Hence, we have

$$\forall t \in T_{x,y}; f(x) \ge f(t) \Rightarrow \exists a \in A_{x,y}; f(a) \ge f(y).$$

We have to show the case with strict inequalities. We prove the contraposition of this: assume $f(a) \leq f(y)$ for all $a \in A_{x,y}$. If $x \in M(S)$, we have $T_{x,y} \cap M(S) \neq \emptyset$, hence there exists some $t \in T_{x,y}$ such that $f(t) \geq f(x)$. Suppose $x \notin M(S)$ and $f(x) \geq f(t)$ for all $t \in T_{x,y}$. Then, there exists $x' \in I_x$ such that $x' \in M(S)$. By our assumption, we must have $T_{x',y} \cap M(S) \neq \emptyset$. However, by Lemma 3, this contradicts the fact that f(x') > f(t). Hence there exists some $t \in T_{x,y}$ such that $f(t) \geq f(x)$. This proves that

$$\forall a \in A_{x,y}; f(a) \le f(y) \Rightarrow \exists t \in T_{x,y}; f(x) \le f(t),$$

which is equivalent to

$$\forall t \in T_{x,y}; f(x) > f(t) \Rightarrow \exists a \in A_{x,y}; f(a) > f(y).$$

(b): Let $x \in M(S)$ and $y \in M(S')$. By s-quasisupermodularity, we have $\forall a \in A_{x,y}; f(a) \geq f(y)$, which means $A_{x,y} \subset M(S')$. Then, suppose there exists some t' such that $t' \notin M(S)$, that is, f(t') < f(x). In this case, by s-quasisupermodularity, we must have f(a) > f(y) for all $a \in A_{x,y}$, contradiction.

The necessity part can be shown as follows. Let S and S' be the same as the proof of (a). Suppose f(y) > f(a) for some $a \in A_{x,y}$. What we have to show is that f(t) > f(x) for all $t \in T_{x,y}$. Note that $I_x \subset (M(S))^c$. This is shown as follows. Suppose $y \in M(S')$ and some $x' \in I_x$ is contained in M(S). In this case, by Lemma 3, we must have $A_{x',y} = A_{x,y} \subset M(S')$, which contradicts our assumption. Suppose some $a' \in A_{x,y}$ is contained in M(S') and some $x' \in I_x$ is an element of M(S). In this case, by Lemmas 2 and 3, we must have $A_{x',a'} = I_{a'} = A_{x,y} \subset M(S')$, which again contradicts our assumption. Hence, the set M(S) consists of some elements of $T_{x,y}$. Let $t' \in T_{x,y}$ be an element of M(S). Note that, in fact, $A_{x,y} \subset (M(S'))^c$. Indeed, if some $a' \in A_{x,y}$ contained in M(S'), then we must have $A_{t',a'} = I_{a'} = A_{x,y} \subset M(S')$, contradiction. Hence, the set M(S') consists of some elements of I_y . Let $y' \in I_y$ be an element of M(S'). Based on the above arguments, we can show that, in fact, $T_{x,y} = M(S)$. Indeed, by Lemmas 2 and 3, we have $T_{t',y'} = I_{t'} = T_{x,y} \subset M(S')$. This proves that

$$\exists a \in A_{x,y}; f(y) > f(a) \Rightarrow \forall t \in T_{x,y}; f(t) > f(x),$$

which is equivalent to

$$\exists t \in T_{x,y}; f(x) \ge f(t) \Rightarrow \forall a \in A_{x,y}; f(a) \ge f(y).$$

The case with strict inequalities is shown as follows: suppose f(x) > f(t) for some $t \in T_{x,y}$. In this case, the set M(S) must consist of some elements of I_x . Let x' be an element of M(S). We show that $I_y \subset (M(S'))^c$. Indeed, if some $y' \in I_y$ is contained in M(S'), then by Lemma 3, we must have $T_{x',y'} = T_{x,y'} = T_{x,y} \subset M(S)$, which contradicts our assumption. Hence, the set M(S') consists of some elements of $A_{x,y}$. Let $a' \in A_{x,y}$ be an element of M(S'). However, by Lemmas 2 and 3, we have $A_{x',a'} = I_{a'} = A_{x,y} \subset M(S')$, which implies $A_{x,y} = M(S')$. This proves that

$$\exists t \in T_{x,y}; f(x) > f(t) \Rightarrow \forall a \in A_{x,y}; f(a) > f(y).$$

This completes our proof. [Q.E.D.]

References

- [1] Milgrom, P., Shannon, C. (1994): Motone comparative statics. *Econometrica*, 62, 157-180.
- [2] Shirai, K. (2008): A generalization of monotone comparative statics. *Economics Bulletin*, Vol. 3, No. 39.